

Identifying Pasture Weeds

Annual poa (*Poa annua*)

Summary Widespread grass weed – abundant in many pastures. Annual.

Identification: Up to 20 cm tall, it grows quickly and heads all year round. It has fine soft light green leaves with tips that curve upwards.

Characteristics: Annual poa grows aggressively in bare areas of pasture in autumn, winter and spring. Typically it finishes seeding and dies in spring when it gets dry. Poa can be abundant in pastures, filling gaps even on compacted ground. Is eaten by stock.



Barley grass (*Critesion* sp.)

Summary: Widespread, particularly in drier close grazed pastures. Annual.

Identification: Has dull green twisted leaves and seed heads that resemble barley.

Characteristics: Does not establish well in dense pasture. Its seed can penetrate the skin of sheep. Seeds are only viable for one to two years so preventing it from seeding can be an effective control method. Few barley grass plants appear following cultivation.



Buttercup (*Ranunculus* sp.)

Summary Creeping buttercup (*Ranunculus repens*) and giant buttercup (*Ranunculus acris*) are the most common. Grow best in wet areas. Perennials.

Identification: Creeping buttercup has runners and grows across the ground, whereas the giant buttercup is more upright. The giant buttercup also has finer more pointed leaves. Both produce yellow flowers.

Characteristics: Eaten by sheep but not cattle. Herbicide resistant. ecotypes of giant buttercup have developed.



specialtyseeds
NZ's INDEPENDENT SEED SUPPLIER

For more detailed product information please ring
0800 727 - 8873 or go to www.specseed.co.nz

Identifying Pasture Weeds

Californian thistle (*Cirsium arvense*)

Summary: Widespread perennial.

Identification Starts growing in November as rosettes from a creeping root system. Leafy stems then develop that can grow 50-150 cm tall. Has small purple flowers that grow from round buds. Seedlings resemble Scotch thistle but grow more slowly and have smoother leaves carrying spines on the edge only.

Characteristics: Typically grow in patches, often from a single plant via its roots.




Chickweed (*Stellaria media*)

Summary: Widespread winter grower that does not like dry conditions. Annual.

Identification A sprawling plant, which has small light green leaves. Produces small white flowers.

Characteristics: Germinates in the autumn and dies during the summer. Can be a problem in autumn sowed pasture because of its ability to grow in shady conditions. Is controlled with frequent grazing through trampling.



Dandelion (*Taraxacum officinale*)

Summary: Widespread perennial.

Identification: Has a taproot and broad leaves with very jagged edges. Produces a bright yellow flower that has a thick hollow unbranched stem.

Characteristics: Its roots have a strong reproductive ability. Pieces of root from as deep as 20 cm can grow into new plants, giving it the capacity to multiply as a result of cultivation. Can thrive in cattle pasture but is kept in check under sheep grazing.



specialtyseeds
NZ's INDEPENDENT SEED SUPPLIER

For more detailed product information please ring
0800 727 - 8873 or go to www.specseed.co.nz

Identifying Pasture Weeds

Dock (broad-leaved) (*Rumex obtusifolius*)

Summary: Widespread - most plentiful on wet soils. Perennial.

Identification: Has large broad leaves and a taproot. Produces buds that turn a reddish colour.

Characteristics: Seeds are long-lived, so large banks of seed can build up in the soil. When dock infested paddocks are cultivated, they are soon reinfested, both from seedlings and regrowth of roots. Sheep eat docks, but cattle and horses avoid them.



Fathen (*Chenopodium album*)

Summary: Spring/summer weed in all districts. Annual.

Identification: The leaves of many seedlings are purple underneath. The first two leaves are horizontal and narrow, but the true leaves are broad. Has an upright central stem. Its seeds ripen and fall while leaves are still green.

Characteristics: Germinates in spring and dies with frosts in autumn. Grows aggressively in dry conditions. Can grow densely in spring sown pasture. Seeds last for 60 years in the soil and can germinate with little rain. Stock can eat it without serious harm but may suffer scouring and ill-thrift.



Nettle (*Urtica urens*)

Summary: Common in the South Island and lower North Island. Annual.

Identification: Recognised by its dark green sharply toothed leaves. When touched, causes stinging for a few minutes from poison released by its fine bristles.

Characteristics: Usually disappears from pasture but can return in patches where pasture is thin. Not eaten by stock.



specialtyseeds
NZ's INDEPENDENT SEED SUPPLIER

For more detailed product information please ring
0800 727 - 8873 or go to www.specseed.co.nz

Identifying Pasture Weeds

Nightshade (black) (*Solanum nigrum*)

Summary: Widespread. Often confused with deadly nightshade. Annual.

Identification: Upright plant that can grow tall and leafy. The leaves and stem can develop a purple tinge, especially late in the season when stressed. It produces a small white flower and bunches of berries that are either green or black.

Characteristics: Germinates in the spring and dies with frosts in autumn. Often confused with deadly nightshade, which is much more poisonous and less common, mainly found around Christchurch. Black nightshade is spread through birds consuming its fruit and spreading seed through their droppings.



Nodding thistle (*Carduus nutans*)

Summary: Common biennial – prefers a dry climate.

Identification: Similar growth habit and size to Scotch thistle. Has red-purple flowers a bit larger than Scotch thistle, and they droop long before maturity.

Characteristics: Is more difficult to control than Scotch thistle. Three insects have been introduced for biological control, one by feeding on the seed in the flower head, one that forms galls and a third that attacks the rosettes. Herbicide-resistant ecotypes have developed in many parts of the country.



Scotch thistle (*Cirsium vulgare*)

Summary: Most widespread thistle. Biennial.

Identification: Has a branched stout stem, which rises in October from a rosette grown the previous season. Plants are short and spindly on poor ground but grow to a height of 1 m in good conditions. Flowers are purple but bluer than nodding thistle.

Characteristics: Scotch thistles often grow after fire or soil disturbance. They can be abundant in pasture as stock, especially sheep, seldom eat them.



specialtyseeds
NZ's INDEPENDENT SEED SUPPLIER

For more detailed product information please ring
0800 727 - 8873 or go to www.specseed.co.nz

Identifying Pasture Weeds

Shepherds purse (*Capsella bursa-pastoris*)

Summary: Common throughout NZ. Annual.

Identification: Its leaves are lobed but vary widely in shape even among plants growing together. It has a central stem (15-50 cm tall) carrying small white flowers and buds that are heartshaped.

Characteristics: It is controlled in pasture by grazing and not letting it seed. The seed is viable for many years in the soil and plants appear in most crops/pasture sowings. It's tolerant of cold conditions and often grows through winter to set seed in spring.



Spurrey (*Spergula arvensis*)

Summary: Widespread annual.

Identification: Has very fine leaves and a sprawling stem. It produces small white flowers, which are gathered at the ends of the stems.

Characteristics: It mostly establishes in the autumn but can establish in spring. Even though it is a small weed it can cause problems by smothering newly sown pastures. Seeds remain viable in the soil for many years and outbreaks can occur following cultivation. Spurrey grows well under acidic soil conditions.



Storksbill (*Erodium cicutarium*)

Summary: Common on the East Coast. Annual to biennial.

Identification: Forms a rosette 10-30 cm in diameter with a deep taproot. Has long narrow pointed fruits.

Characteristics: Germinates in autumn and summer. In pasture, this weed provides reasonable feed in dry conditions but the fruits can penetrate the skin of sheep.



specialtyseeds
NZ's INDEPENDENT SEED SUPPLIER

For more detailed product information please ring
0800 727 - 8873 or go to www.specseed.co.nz

Identifying Pasture Weeds

Wireweed (*Polygonum aviculare*)

Summary: Widespread annual.

Identification: Has a prostrate growth habit. Its stems are tough and branch up to 150 cm long. Small leaves are always green and seedlings have semi-erect leaves. It produces small flowers at the base of the leaves that are often hard to notice.

Characteristics: Closely related to willow weed. Establishes in spring, earlier than most summer annuals. Does not persist in a dense pasture sward. Needs good moisture for germination but is tolerant of low temperatures. Very tolerant of treading.



specialtyseeds
NZ's INDEPENDENT SEED SUPPLIER

For more detailed product information please ring
0800 727 - 8873 or go to www.specseed.co.nz